

Copper Fungicide

(Stone Fruit: Almond, Apricot, Cherry, Nectarine, Peach, Plum)

1. Bacterial Canker

Apply as a dormant spray at bud swell and at bud burst stage, weekly thereafter.

2. Brown Rot Blossom

Apply at bud swell, popcorn, full bloom, and petal fall.

3. Pome Fruit- (Apple, Pear, Quince)

Fruit Tree Spray

(Apple, Cherry, Grape, Peach) • Per Gallon rates / 10-14 day intervals*

1. Apple- black/white rots - botrytis rot - brown rot blossom - cedar rust - scab

1.5 tbs* at green tip / 2.5 tbs* at pre bloom, full pink, petal fall to 7-10 days after petal fall

2. Cherry- botrytis rot - brown rot blossom blight - fruit rot - cherry leaf spot

1.5 tbs* at green tip / 2.5 tbs* at pre bloom, full pink, petal fall to 7-10 days after petal fall

3. Grape- black rot - botrytis rot

1.5 tbs* at green tip / 2.5 tbs* at pre bloom, full pink, petal fall to 7-10 days after petal fall

4. Peach- brown rot (fruit) - rhizopus - scab - coryneum blight

1.5 tbs* at green tip / 1.5 tbs* at pre bloom / 2.5 tbs* at full pink / 2.5 tbs* 7-10 days after petal fall

Fung-onil Multi Purpose Fungicide (Peach, Nectarine, Apricot, Cherry, Plum, Prune)

Do not apply within one week before or after application of oil.

1. Leaf Curl

3.75 tsp in 4 gallons of water applied at leaf fall in Autumn and 1-2 times before bud swell

2. Shothole, brown rot blossom blight, cherry leaf spot

3.75 tsp in 4 gallons of water at bud break or pop corn. Second application 10 days later (full bloom to petal fall). Apply at shuck split to prevent infections on young fruit.

3. Blueberry - Mummy Berry

3.5 tsp in 4 gallons of water. Apply at green tip. Repeat applications until early bloom at 10 day intervals. Do not apply after early bloom.

Sulphur Plant Fungicide (Apples, Blackberries, Blueberries, Boysenberries, Huckleberries, Loganberries, Cherries, Grapes, Nectarines, Peaches, Plums, Prunes, Pears, Roses)

1. Berries

(powdery mildew) 2-2.5 tbs per gallon. Begin when new shoots are 6 inches long and before blossoms open.

2. Apples

(apple cedar rust, leaf spot, black/brown rot, scab) 1-2.5 tbs per gallon from pre pink through 7-10 days after petal fall.

3. Grapes

(powdery mildew) 1-3 tbs per gallon. Begin when shoots are 6 inches long. Repeat before blossoms open.

4. Nectarines & Peaches

(brown rot, peach canker, rust, scab, shothole) 1-3 tbs per gallon. Apply at pink, bloom, petal fall, shuck split, shuck fall.

All recommendations are excerpts from the label of products, consult labels for full instruction and application.